eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7

# PAP SMEAR SCREENING: INSIGHTS INTO KNOWLEDGE, ATTITUDE, AND PRACTICE AMONG WOMEN IN ABU DHABI'S PRIMARY HEALTHCARE CENTERS

**Nasir Shaima** 

Sheikh Khalifa Medical City, P.O.Box: 51900, Abu Dhabi, United Arab Emirates

#### **ABSTRACT**

This study investigates the knowledge, attitude, and practice (KAP) of Pap smear screening among women attending primary healthcare centers in Abu Dhabi Island. Pap smear screening is a vital tool for early detection of cervical cancer, yet its utilization is influenced by various factors including awareness, perception, and accessibility. Through a structured survey, data was collected from women visiting primary healthcare centers, focusing on their understanding of Pap smear screening, attitudes towards the procedure, and actual practice of undergoing screenings. The findings shed light on gaps in knowledge, misconceptions, and barriers to Pap smear utilization, offering insights for targeted interventions to improve cervical cancer screening rates and women's health outcomes in Abu Dhabi.

#### **KEYWORDS**

Pap smear screening, Cervical cancer, Knowledge, Attitude, Practice, Women's health, Primary healthcare centers, Abu Dhabi.

## **INTRODUCTION**

Cervical cancer is a significant global health concern and remains one of the leading causes of cancer-related mortality among women worldwide. Despite advances in prevention and treatment, cervical cancer continues to pose a substantial burden on women's health, particularly in regions with limited access to screening and healthcare services. Abu Dhabi, the capital city of the United Arab Emirates (UAE), is no exception to this public health challenge.

Pap smear screening, a simple and effective test for detecting precancerous and cancerous changes in the cervix, plays a crucial role in early detection and prevention of cervical cancer. However, the utilization of Pap smear screening services is influenced by various factors, including women's knowledge about the procedure, their attitudes towards screening, and their actual practice of undergoing screenings.

This study aims to investigate the knowledge, attitude, and practice (KAP) of Pap smear screening among women attending primary healthcare centers in Abu Dhabi's island region. By examining women's understanding of Pap smear screening, their perceptions of the procedure, and their actual utilization of screening services, this research seeks to identify barriers and facilitators to cervical cancer screening in Abu

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7

## Dhabi.

The findings of this study are expected to provide valuable insights into the gaps in knowledge, misconceptions, and barriers that hinder women's access to Pap smear screening services in Abu Dhabi. Understanding these factors is critical for developing targeted interventions and strategies to improve cervical cancer screening rates and promote women's health in the region.

Through a structured survey administered to women visiting primary healthcare centers, this study will assess the level of awareness and understanding of Pap smear screening, explore women's attitudes towards the procedure, and examine the factors influencing their decision to undergo screening. By elucidating the KAP of Pap smear screening among women in Abu Dhabi, this research aims to contribute to the development of evidence-based interventions to enhance cervical cancer prevention efforts and improve women's health outcomes in the region.

In the subsequent sections, we will delve into the methodology employed in this study, including the survey design, data collection procedures, and analysis methods. We will also discuss the theoretical framework guiding this research and the significance of understanding Pap smear screening KAP in the context of cervical cancer prevention and women's health promotion in Abu Dhabi's primary healthcare centers.

### **METHOD**

The process of examining Pap smear screening among women in Abu Dhabi's primary healthcare centers involved a systematic and structured approach. Initially, the research team collaborated with healthcare authorities to gain access to primary healthcare centers across Abu Dhabi Island. Through strategic planning, the team employed a stratified random sampling technique to ensure representation from diverse demographic backgrounds among the participants.


The development of the survey instrument was a critical step in the process. The research team designed a structured questionnaire to capture information on participants' knowledge, attitude, and practice regarding Pap smear screening. The questionnaire underwent a rigorous validation process, including expert review and pilot testing, to ensure clarity, comprehensibility, and relevance to the study objectives.

Data collection was carried out by trained research assistants in a professional and sensitive manner. Participants were approached in primary healthcare centers and provided with detailed information about the study objectives and procedures. Informed consent was obtained from all participants before administering the questionnaire. The research team ensured privacy and confidentiality throughout the data collection process, emphasizing the voluntary nature of participation and the anonymity of responses.

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7



Following data collection, the research team meticulously organized and analyzed the data using appropriate statistical methods and software packages. Descriptive statistics were utilized to summarize participants' demographic characteristics, knowledge levels, attitudes, and screening practices. Inferential statistics, including chi-square tests and logistic regression analysis, were employed to explore associations between demographic variables and Pap smear screening outcomes.

Ethical considerations were paramount throughout the research process. The study obtained ethical approval from the Institutional Review Board (IRB) to ensure compliance with ethical guidelines and standards for research involving human participants. Participants' rights to privacy, confidentiality, and informed consent were rigorously upheld, reflecting the ethical principles underpinning the study.

While acknowledging the limitations inherent in the study design, including potential biases and constraints associated with self-reported data, the research team strived to mitigate these limitations through rigorous methodological procedures and data validation techniques. The insights generated from this study have the potential to inform healthcare policies, interventions, and educational campaigns aimed at enhancing Pap smear screening rates and promoting women's health in Abu Dhabi's primary healthcare centers.

A stratified random sampling technique was employed to recruit participants from primary healthcare centers


eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7

across Abu Dhabi Island. Women aged 21 and above, who were attending primary healthcare centers for routine check-ups or consultations, were invited to participate in the study. Efforts were made to ensure representation from diverse demographic backgrounds and socioeconomic statuses.

A structured questionnaire was developed to assess participants' knowledge, attitude, and practice (KAP) regarding Pap smear screening. The questionnaire included sections on demographic information, knowledge about cervical cancer and Pap smear screening, attitudes towards screening, barriers to screening, and past screening practices. The questionnaire was reviewed by experts in women's health and validated through pilot testing with a small sample of participants.



Trained research assistants administered the questionnaire to eligible participants in private consultation rooms at primary healthcare centers. Participants were briefed about the purpose of the study and provided with information about the voluntary nature of their participation. Informed consent was obtained from all participants before proceeding with data collection. Confidentiality and anonymity of participants' responses were ensured throughout the study.

Quantitative data collected through the survey were entered into a statistical software package for analysis. Descriptive statistics, including frequencies, percentages, means, and standard deviations, were calculated to summarize participants' demographic characteristics, knowledge scores, attitude scores, and screening


eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7

practices. Inferential statistics, such as chi-square tests and logistic regression analysis, were used to explore associations between demographic variables and Pap smear screening outcomes.

Ethical approval for the study was obtained from the Institutional Review Board (IRB) of the relevant healthcare authority in Abu Dhabi. The study adhered to ethical guidelines for research involving human participants, including informed consent, privacy protection, and confidentiality of data. Participants were assured of their right to withdraw from the study at any time without consequence.



Several limitations of the study should be acknowledged, including the potential for selection bias due to the convenience sampling method and the reliance on self-reported data, which may be subject to recall bias and social desirability bias. Despite these limitations, efforts were made to minimize biases and enhance the validity and reliability of findings through rigorous survey design and data collection procedures.

By employing a structured survey instrument and systematic sampling approach, this study aimed to provide a comprehensive understanding of women's knowledge, attitude, and practice regarding Pap smear screening in Abu Dhabi's primary healthcare centers. The insights generated from this study have the potential to inform targeted interventions and policies aimed at improving cervical cancer screening rates and promoting women's health in the region.

### **RESULT**

The study on Pap smear screening among women in Abu Dhabi's primary healthcare centers revealed several

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7

key findings regarding knowledge, attitude, and practice (KAP) related to cervical cancer screening. Firstly, the majority of participants demonstrated limited knowledge about cervical cancer and Pap smear screening, with misconceptions and gaps in understanding prevalent among the respondents. Despite the availability of screening services in primary healthcare centers, many women lacked awareness of the importance of regular screenings for early detection and prevention of cervical cancer.

Attitudinally, while participants generally expressed positive attitudes towards Pap smear screening and recognized its importance in detecting cervical abnormalities, concerns and misconceptions about the procedure persisted. Fear of pain, embarrassment, and cultural barriers were commonly cited factors influencing women's attitudes towards undergoing Pap smear screening.

In terms of practice, the study revealed suboptimal utilization of Pap smear screening services among women attending primary healthcare centers in Abu Dhabi. Many participants reported never having undergone a Pap smear test, citing reasons such as lack of awareness, perceived invasiveness of the procedure, and cultural norms surrounding women's health.

### **DISCUSSION**

The findings underscore the urgent need for targeted interventions and educational campaigns to enhance knowledge, shift attitudes, and promote uptake of Pap smear screening among women in Abu Dhabi. Addressing misconceptions, alleviating fears, and raising awareness about the importance of cervical cancer screening are crucial steps towards improving women's health outcomes and reducing the burden of cervical cancer in the region.

Cultural sensitivity and community engagement are paramount in designing effective interventions aimed at increasing Pap smear screening rates. Culturally tailored educational materials, outreach programs, and public health initiatives can help dispel myths, normalize discussions about women's health, and empower women to make informed decisions about their reproductive health.

Healthcare providers also play a pivotal role in promoting Pap smear screening and facilitating access to screening services. Patient-centered approaches, such as counseling, education, and support, can help address women's concerns and encourage uptake of screening services in primary healthcare settings.

## CONCLUSION

In conclusion, the study highlights the need for comprehensive strategies to improve knowledge, shift attitudes, and promote uptake of Pap smear screening among women in Abu Dhabi's primary healthcare centers. By addressing barriers, dispelling myths, and raising awareness about the importance of cervical cancer screening, healthcare authorities can enhance women's access to preventive services and contribute to reducing the incidence and mortality associated with cervical cancer in the region.

Moving forward, collaborative efforts between healthcare providers, policymakers, community organizations, and advocacy groups are essential in driving meaningful change and improving women's health outcomes through increased Pap smear screening uptake. By prioritizing women's health and investing in preventive care initiatives, Abu Dhabi can take significant strides towards achieving better cervical cancer outcomes and promoting overall well-being among its population.

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE05 Published 02-05-2023

Page No. 1-7

## **REFERENCES**

- **1.** Health authority Abu-Dhabi website. Wave Health Promotion Project 2012-2013. http://www.haad.ae/haad/tabid/1257/Default.aspx (accessed 3rd February 2013).
- **2.** WHO. WHO cancer fact sheet. http://www.who.int/media-centre/factsheets/fs297/en/index.html (accessed 3rd February 2013).
- 3. Bakheit, N.M. and BuHaroon, A.I. THE KNOWLEDGE, ATTITUDE AND PRACTICE OF PAP-SMEAR AMONG LOCAL SCHOOL TEACHERS IN THE SHARJAH DISTRICT. Middle East Journal of Family Medicine.2004;4(): 45
- **4.** Middle east health magazine. HPV prevalence in the Middle East. http://www.middleeasthealthmag.com/cgibin/index.cgi?http://www.middleeasthealthmag.com/may2 011/feature9.htm (accessed 3rd February 2013).
- 5. Nakalevu, S.M. e Knowledge, Attitude, Practice and Behavior of Women Towards Cervical Cancer and Papsmear Screening A cross sectional study in the Rewa subdivision. http://www.pacifichealthvoices.org/files/Knowledge,%20Attitude\_to%20 Cervical%20Ca.pdf (accessed 3rd February 2013).
- **6.** Gharoro, E.P., And Ikeanyi E.N. An appraisal of the level of awareness and utilization of the Pap-smear as a cervical cancer screening test among female health workers in a tertiary health institution. International Journal of Gynecology Cancer .2006;16(16):1063-1068
- 7. Health authority Abu-Dhabiwebsite. HAAD Standards for the Cervical Cancer Screening. http://www.haad.ae/HAAD/Link-Click.aspx?fileticket=M\_AHzOF-aRw%3d&tabid=819 (accessed 3rd February 2013).
- **8.** Noureddine,M., Kabakian, S. A survey of knowledge, attitude, and practice of cervical screening among Lebanese / American women. nursing outlook.1997;45(1):16-22
- **9.** Al-Meer,F.M., Aseel,M.T.,Al-Khalaf, J., Al-Kuwari,M.G. and Ismail,M.F.S. 'Knowledge, attitude and practices regarding cervical cancer and screening among women visiting primary health care in Qatar. eastern mediterrenean health journal.2011;17(11):855-861
- **10.** Gamarra C.J, Paz E.P, Griep R.H. 'Knowledge, attitudes and practice related to Papanicolaou smear test among Argentina's women. .2005;39(2):270