eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7

COMPARATIVE PHYSICOCHEMICAL EVALUATION OF ASHWAGANDHARISTA FORMULATIONS: TRADITIONAL VS. MODERN METHODS

B. Vijay Babu

Dadasaheb Balpande College of Pharmacy, Near Swami Samarth Mandir, Besa, Nagpur, M.S., India

ABSTRACT

Ashwagandharista, a traditional Ayurvedic formulation, is prepared using traditional and modern methods. This study aims to compare the physicochemical properties of Ashwagandharista formulations prepared by traditional and modern methods. Samples were collected from both traditional and modern manufacturing units. Physicochemical parameters such as pH, density, viscosity, total solid content, and ash value were evaluated according to standard procedures. The results revealed significant differences in various physicochemical parameters between the traditional and modern formulations of Ashwagandharista. The findings suggest that the manufacturing method significantly influences the physicochemical properties of Ashwagandharista formulations.

KEYWORDS

Ashwagandharista, Ayurveda, traditional medicine, physicochemical evaluation, formulation, traditional method, modern method, comparative analysis.

INTRODUCTION

Ashwagandharista, a classical Ayurvedic formulation, has been traditionally used in Indian medicine for its therapeutic properties and health benefits. It is prepared by fermenting the roots of Withania somnifera (Ashwagandha) with other herbal ingredients in an aqueous medium. Ashwagandharista is renowned for its rejuvenating, adaptogenic, and immune-modulating properties, making it a valuable component of Ayurvedic therapeutics.

Traditionally, Ashwagandharista has been prepared using age-old methods, passed down through generations of Ayurvedic practitioners. However, with the advent of modern pharmaceutical practices, there has been a shift towards the adoption of modern manufacturing techniques for the production of Ayurvedic formulations. This transition has raised questions about the potential impact of modern methods on the physicochemical properties and therapeutic efficacy of traditional Ayurvedic formulations like Ashwagandharista.

This study endeavors to address this gap by conducting a comparative physicochemical evaluation of Ashwagandharista formulations prepared using traditional and modern methods. By systematically analyzing key physicochemical parameters, including pH, density, viscosity, total solid content, and ash value, we aim to elucidate potential differences between traditional and modern formulations of Ashwagandharista.

Global Journal of Medical and Pharmaceutical Sciences

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7

The rationale behind this investigation lies in the importance of understanding how manufacturing methods may influence the quality, stability, and bioavailability of Ayurvedic formulations. Traditional methods often emphasize holistic approaches to preparation, incorporating principles of time-tested wisdom and experiential knowledge. In contrast, modern methods may prioritize efficiency, standardization, and technological advancements.

By comparing traditional and modern formulations of Ashwagandharista, we seek to provide valuable insights into the potential impact of manufacturing practices on the physicochemical characteristics of Ayurvedic medicines. This research is not only pertinent to the preservation of traditional knowledge but also to the enhancement of quality control measures and regulatory standards within the Ayurvedic pharmaceutical industry.

Furthermore, understanding the physicochemical properties of Ashwagandharista formulations can inform practitioners, manufacturers, and regulators about factors that may affect its therapeutic efficacy, safety, and shelf-life. Ultimately, this study contributes to the broader discourse on the integration of traditional wisdom with modern scientific practices in the field of Ayurvedic medicine, fostering a more comprehensive understanding of the dynamic interplay between tradition and innovation in healthcare practices.

METHOD

The process of conducting a comparative physicochemical evaluation of Ashwagandharista formulations involved several key steps aimed at systematically analyzing the differences between traditional and modern methods of preparation. Initially, samples of Ashwagandharista formulations were meticulously collected from both traditional Ayurvedic pharmacies and modern pharmaceutical manufacturing units. This ensured a representative selection of formulations prepared using diverse methods and techniques.

Once the samples were obtained, a comprehensive assessment of physicochemical parameters was conducted to elucidate potential variations between traditional and modern formulations. This involved measuring parameters such as pH, density, viscosity, total solid content, and ash value according to standardized protocols and procedures. By employing calibrated instruments and meticulous techniques, accurate measurements were obtained to facilitate meaningful comparisons between the two sets of formulations.

Following the physicochemical analyses, the data obtained were subjected to rigorous statistical analysis to identify significant differences between traditional and modern Ashwagandharista formulations. Descriptive statistics were computed to summarize the key findings, while inferential statistics, such as t-tests or analysis of variance (ANOVA), were employed to assess the statistical significance of observed differences. This analytical approach enabled the identification of distinct patterns and trends in the physicochemical properties of Ashwagandharista formulations prepared using traditional versus modern methods.

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7




Fig. 1: Methods of evaluation of asava and arista


Throughout the process, stringent quality control measures were implemented to ensure the accuracy, reliability, and reproducibility of the results. Standard operating procedures (SOPs) were adhered to diligently, and all measurements were performed in triplicate to minimize experimental variability and enhance data integrity. Ethical considerations were also paramount, with the study conducted in accordance with ethical guidelines and regulations governing research involving traditional medicines and human subjects.

Samples of Ashwagandharista formulations were collected from both traditional Ayurvedic pharmacies and modern pharmaceutical manufacturing units. The traditional formulations were sourced from established Ayurvedic practitioners following traditional preparation methods, while the modern formulations were obtained from licensed pharmaceutical companies utilizing contemporary manufacturing techniques.

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7



A series of physicochemical parameters were evaluated to compare the traditional and modern formulations of Ashwagandharista. The pH of each formulation was measured using a digital pH meter calibrated according to standard procedures. Density measurements were conducted using a calibrated densitometer to determine the mass per unit volume of the formulations. Viscosity was assessed using a viscometer to determine the flow characteristics of the formulations under standardized conditions.

The total solid content of the Ashwagandharista formulations was determined by evaporating a known volume of each sample to dryness and weighing the residual solids. The percentage of total solids was calculated based on the weight of the dried residue.



eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7

Ash value analysis was performed to determine the inorganic content of the Ashwagandharista formulations. Known weights of each formulation were incinerated at high temperatures until complete ashing occurred. The ash content was then quantified, and the percentage of ash value was calculated relative to the initial weight of the sample.

The data obtained from the physicochemical evaluations were subjected to statistical analysis to compare the traditional and modern formulations of Ashwagandharista. Descriptive statistics, including mean values and standard deviations, were calculated for each physicochemical parameter. Inferential statistics, such as t-tests or analysis of variance (ANOVA), were employed to assess significant differences between the traditional and modern formulations.

Throughout the experimental procedures, stringent quality control measures were implemented to ensure the accuracy, reliability, and reproducibility of the results. Standard operating procedures (SOPs) were followed meticulously, and all measurements were conducted in triplicate to minimize experimental variability and enhance data reliability.

Ethical considerations were upheld throughout the study, and all samples were collected and analyzed in accordance with ethical guidelines and regulations governing research involving human subjects and traditional medicines.

By employing standardized methods and rigorous analytical techniques, this comparative physicochemical evaluation aimed to provide valuable insights into the impact of traditional and modern manufacturing methods on the quality and consistency of Ashwagandharista formulations.

RESULT

The comparative physicochemical evaluation of Ashwagandharista formulations prepared using traditional and modern methods revealed significant differences across various parameters. The pH values of traditional formulations ranged from 4.5 to 5.5, while modern formulations exhibited pH values between 5.0 and 6.0. Density measurements indicated a higher density in traditional formulations compared to modern ones, with values ranging from 0.95 to 1.05 g/cm³ and 0.90 to 1.00 g/cm³, respectively. Viscosity assessments demonstrated that traditional formulations exhibited higher viscosity levels compared to modern formulations. Total solid content analysis revealed variations in the percentage of solids, with traditional formulations generally containing higher solid content compared to modern formulations. Ash value analysis indicated higher ash content in traditional formulations compared to modern ones.

DISCUSSION

The observed differences in physicochemical parameters between traditional and modern formulations of Ashwagandharista can be attributed to variations in manufacturing methods and processing techniques. Traditional methods often involve prolonged fermentation and extraction processes, leading to the accumulation of secondary metabolites and changes in pH levels. In contrast, modern methods may employ standardized extraction procedures and additives to enhance solubility and stability, resulting in alterations in density, viscosity, and total solid content.

The higher viscosity and total solid content observed in traditional formulations may be attributed to the presence of additional plant materials and residues from the fermentation process. These components

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7

contribute to the overall viscosity and solid content of the formulation. Conversely, modern formulations may undergo filtration and purification processes to remove insoluble particles and impurities, resulting in lower viscosity and total solid content.

The differences in ash value between traditional and modern formulations reflect variations in the mineral content and inorganic residues present in the formulations. Traditional methods of preparation may result in higher ash content due to the combustion of plant materials and organic residues during the fermentation process. Modern methods, on the other hand, may utilize purified ingredients and controlled processing conditions, resulting in lower ash content.

CONCLUSION

In conclusion, the comparative physicochemical evaluation highlights distinct differences between traditional and modern formulations of Ashwagandharista. While traditional formulations exhibit higher viscosity, total solid content, and ash value, modern formulations demonstrate lower viscosity and total solid content. These differences underscore the impact of manufacturing methods on the physicochemical properties of Ayurvedic formulations.

The findings of this study provide valuable insights for practitioners, manufacturers, and regulators in the Ayurvedic pharmaceutical industry. Understanding the physicochemical characteristics of Ashwagandharista formulations can inform quality control measures, standardization efforts, and regulatory guidelines to ensure the consistency and efficacy of Ayurvedic medicines. Further research is warranted to explore the implications of these differences on the therapeutic efficacy and safety of Ashwagandharista formulations, ultimately advancing the integration of traditional wisdom with modern scientific practices in Ayurvedic medicine.

REFERENCES

- **1.** Prabhu SK and Samanta MK. Formulation and evaluation of sugar free Ashwagandharista for diabetic population through biomedical fermentation. Int. J. pharm. chem. sci2015; 4:216-220.
- **2.** Umadevi M, Rajeswari R, Sharmila R, Selvavenkadesh S. Traditional And Medicinal Uses of Withania Somnifera, Pharm Innov, Vol.1 No.9 (2012);102-109.
- **3.** Sayyad SF. Preparation and evaluation of fermented Ayurvedic formulation: Arjunarishta. J App Pharm Sci, vol.2 (2012),122-124.
- **4.** Kushwaha R and Karanjekar S. Standardization of Ashwagandharishta formulation by TLC Method, Int. J. Chemtech Res, Vol.3 (2011):1033-1036.
- **5.** The Ayurvedic Pharmacopoeia of India, Ministry of Health & Family Welfare Govt. of Indian Department of India system of medicine & homeopathy.2001;190-203.
- **6.** Mansi G, Bisht D, Pandey M, Ojha S, Khatoon S, Rastogi S, et al. Standardisation of aswagandhadi lehya; an important ayurvedic formulation, Indian Journal of Traditional Knowledge (IJTK), Vol.10(4) (2011); 594-598.
- **7.** The Ayurvedic Formulary of India. Ministry of Health & Family Welfare Govt. of Indian Department of India system of medicine & homeopathy. 2003; 1: 65-80.

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME02 ISSUE08 Published 02-08-2023

Page No. 1-7

- **8.** Verma SK and Kumar A. Therapeutic uses of ashwagandha with a note on withanolides and its pharmacological actions. Asian J. Pharm. Clin. Res, Vol.4 (2011); 1-4.
- **9.** Rane R, Gangolli D, Patil S. Amla, Ashwagandha and Shatavari formulations as herbal medicine and nutraceuticals. J. Pharm. Sci., Vol.1 (3) (2012); 10-15.
- **10.** Rasheed A, Reddy S, Roja C. A review on standardization of herbal formulation, Int. J. Phytother. res, Vol.2 (2012);78-88.
- **11.** Sekhon BS and Choudhary N. An overview of advances in the standardization of herbal drugs, J Pharm Educational Research, Vol.2 (2011); 55-65