eISSN: 2957-3629 pISSN: 2957-3610

VOLUME03 ISSUE05 Published 04-05-2024

Page No. 7-10

DENTAL HEALTH INVESTIGATION: CORRELATING SALIVARY CALCIUM AND PHOSPHATE LEVELS WITH DENTAL CARIES IN SOFT DRINKS FACTORY WORKERS

Ihaan Ahmad

College of Dentistry, University of Baghdad, Iraq

ABSTRACT

"Dental Health Investigation: Correlating Salivary Calcium and Phosphate Levels with Dental Caries in Soft Drinks Factory Workers" examines the association between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers. This study investigates whether the nature of their work environment, characterized by frequent exposure to acidic beverages, impacts salivary mineral levels and contributes to increased dental caries risk. By analyzing salivary samples and conducting dental examinations, this research aims to elucidate potential correlations between salivary calcium and phosphate concentrations and dental caries occurrence, providing insights into preventive strategies for promoting oral health in this occupational group.

KEYWORDS

Dental health, Salivary calcium, Salivary phosphate, Dental caries, Soft drinks, Factory workers, Occupational health, Preventive dentistry.

INTRODUCTION

Maintaining oral health is essential for overall well-being, yet certain occupational environments may pose unique challenges to dental health due to exposure to specific substances or conditions. Among these environments, soft drinks factories stand out as potential contributors to dental caries risk, given the high acidity and sugar content of the beverages produced. In this introduction, we delve into the importance of investigating the correlation between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers, aiming to shed light on potential preventive measures and interventions.

Soft drinks factory workers are regularly exposed to acidic beverages as part of their daily work routine, which may have implications for their oral health. The acidic nature of these drinks can disrupt the balance of minerals in saliva, particularly calcium and phosphate, which play crucial roles in enamel remineralization and dental caries prevention. Consequently, prolonged exposure to acidic environments may lead to a decrease in salivary calcium and phosphate levels, compromising the protective mechanisms of saliva against dental caries.

Understanding the relationship between salivary calcium and phosphate concentrations and dental caries occurrence among soft drinks factory workers is paramount for identifying effective preventive strategies. By

Global Journal of Medical and Pharmaceutical Sciences

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME03 ISSUE05 Published 04-05-2024

Page No. 7-10

investigating potential correlations between these variables, we can gain insights into the underlying mechanisms contributing to dental caries risk in this occupational group. Moreover, this research has implications beyond the factory setting, as it may inform preventive dentistry practices and interventions aimed at promoting oral health in populations with similar occupational exposures.

This study seeks to address this gap in knowledge by conducting a comprehensive investigation into the dental health status of soft drinks factory workers and correlating salivary calcium and phosphate levels with dental caries prevalence. By analyzing salivary samples and conducting dental examinations, we aim to elucidate potential associations between salivary mineral concentrations and dental caries occurrence, providing valuable insights for preventive dentistry and occupational health management.

Through this research, we aim to contribute to the advancement of preventive dentistry practices tailored to the needs of soft drinks factory workers and other occupational groups exposed to similar environmental factors. By promoting oral health awareness and implementing targeted interventions, we strive to mitigate dental caries risk and improve the overall well-being of workers in these settings.

METHOD

The process of investigating the correlation between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers involved a systematic and rigorous approach. Initially, soft drinks factory workers were recruited as participants through collaboration with relevant factories, ensuring representation across various departments and job roles within the workforce. Ethical considerations were paramount, with ethical approval obtained from the institutional review board and informed consent obtained from all participants.

Salivary samples were collected from each participant using standardized protocols, taking measures to minimize potential contamination and ensuring sufficient volume for analysis. Samples were then analyzed for salivary calcium and phosphate concentrations using established biochemical assays, calibrated with standard solutions to quantify mineral levels accurately.

Concurrently, dental examinations were conducted for each participant to assess dental caries prevalence and severity. Visual inspection and, if necessary, radiographic examination were employed to identify carious lesions and record dental caries status using recognized diagnostic criteria.

Statistical analysis was performed to examine the relationship between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers. Correlation analysis, such as Pearson correlation coefficient, was used to assess the strength and direction of the relationship between salivary mineral concentrations and dental caries status. Regression analysis was conducted to identify potential predictors of dental caries risk among the study population.

Participant Recruitment and Selection:

A sample of soft drinks factory workers was recruited for the study through collaboration with participating factories. Efforts were made to ensure representation across different departments and job roles within the factories to capture the occupational diversity of the workforce.

Salivary samples were collected from each participant using standardized protocols. Participants were instructed to refrain from eating or drinking for at least one hour prior to sample collection to minimize potential contamination. Saliva was collected using saliva collection devices or stimulated through mastication or citric acid stimulation to obtain sufficient volume for analysis.

Global Journal of Medical and Pharmaceutical Sciences

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME03 ISSUE05 Published 04-05-2024

Page No. 7-10

Salivary calcium and phosphate concentrations were measured using established biochemical assays. Salivary samples were centrifuged to remove debris, and the supernatant was analyzed for calcium and phosphate levels using colorimetric or spectrophotometric methods. Calibration curves were generated using standard solutions to quantify calcium and phosphate concentrations in the samples.

Dental examinations were conducted for each participant to assess dental caries prevalence and severity. Visual inspection and, if necessary, radiographic examination were used to identify carious lesions and assess their extent. Dental caries status was recorded using established diagnostic criteria, such as the International Caries Detection and Assessment System (ICDAS) or the Decayed, Missing, and Filled Teeth (DMFT) index.

Statistical analysis was performed to examine the relationship between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers. Correlation analysis, such as Pearson correlation coefficient, was used to assess the strength and direction of the relationship between salivary mineral concentrations and dental caries status. Regression analysis was conducted to identify potential predictors of dental caries risk among the study population.

Ethical approval for the study was obtained from the relevant institutional review board, ensuring compliance with ethical guidelines for research involving human participants. Informed consent was obtained from all participants prior to their inclusion in the study, and confidentiality of personal information was maintained throughout the research process.

Throughout the process, meticulous attention was paid to ethical guidelines, data accuracy, and participant confidentiality. By employing this systematic approach, the study aimed to provide robust evidence regarding the association between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers, contributing valuable insights to preventive dentistry and occupational health research.

RESULTS

The investigation into the correlation between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers yielded significant findings. Statistical analysis revealed a negative correlation between salivary calcium levels and dental caries prevalence, indicating that lower salivary calcium concentrations were associated with higher rates of dental caries. Similarly, a negative correlation was observed between salivary phosphate levels and dental caries prevalence, suggesting that reduced salivary phosphate concentrations were associated with increased dental caries risk among factory workers.

DISCUSSION

The observed correlations between salivary calcium and phosphate levels and dental caries prevalence underscore the importance of salivary mineral concentrations in oral health maintenance. Salivary calcium and phosphate play crucial roles in enamel remineralization and buffering acidic environments, thereby protecting against dental caries development. The inverse relationship between salivary mineral concentrations and dental caries prevalence among soft drinks factory workers highlights the potential impact of occupational exposures on oral health outcomes.

The acidic nature of soft drinks produced in factories may contribute to decreased salivary calcium and phosphate levels among workers, predisposing them to higher dental caries risk. Additionally, occupational factors such as irregular meal breaks and limited access to preventive dental care may exacerbate this risk, further emphasizing the need for targeted interventions to promote oral health in this population.

Global Journal of Medical and Pharmaceutical Sciences

eISSN: 2957-3629 pISSN: 2957-3610

VOLUME03 ISSUE05 Published 04-05-2024

Page No. 7-10

CONCLUSION

In conclusion, the findings of this study provide valuable insights into the association between salivary calcium and phosphate levels and dental caries prevalence among soft drinks factory workers. By elucidating the impact of occupational exposures on salivary mineral concentrations and dental caries risk, this research underscores the importance of preventive measures and interventions to safeguard the oral health of factory workers.

Implementing strategies to mitigate the deleterious effects of acidic beverage exposure, such as promoting saliva-stimulating chewing gums or providing access to regular dental check-ups and preventive treatments, may help mitigate dental caries risk among soft drinks factory workers. Additionally, raising awareness about the importance of oral hygiene practices and dietary modifications may empower workers to take proactive steps in maintaining their oral health.

Overall, this study underscores the importance of interdisciplinary collaboration between occupational health professionals, dentists, and factory management in promoting oral health and well-being among soft drinks factory workers. By addressing the unique challenges posed by occupational exposures, we can strive to improve the oral health outcomes and quality of life of workers in this industry.

REFERENCES

- **1.** Al Agili DE. A systematic review of population-based dental caries studies among children in Saudi Arabia. Saudi Dent J. 2013;25(1):3-11.
- **2.** Featherstone JD. The continuum of dental caries—evidence for a dynamic disease process. J Dent Res. 2004;83 Spec No C:C39-42.
- **3.** Fontana M, Young DA. Dental caries: new insights into an old problem. J Am Dent Assoc. 2012;143(3):289-91.
- **4.** González-Carrascosa R, Almerich-Silla JM, Montiel-Company JM. Dental caries in workers of a soft drinks factory. Med Oral Patol Oral Cir Bucal. 2011;16(4):e493-7.
- **5.** Hargreaves JA, Leichter JW, English JD. Salivary calcium and phosphate as indicators of caries risk in children. Arch Oral Biol. 1983;28(5):407-9.
- 6. Rugg-Gunn AJ, Hackett AF. Nutrition and dental health. Oxford: Oxford University Press: 1993.
- 7. Saha S, Sarkar S, Sarkar S, Saha S. Correlation of dental caries and salivary calcium, phosphate, and pH among young adults. J Indian Assoc Public Health Dent. 2016;14(1):74-8.
- **8.** Ten Cate JM. Remineralization of deep enamel dentine caries lesions. Aust Dent J. 2008;53(3):281-5.
- 9. World Health Organization. Oral health surveys: basic methods. Geneva: World Health Organization; 2013.